

Joint statement  
on the role  
of respiratory rehabilitation  
in the COVID-19 crisis:  
the Italian position paper

## *A cura di*

Michele Vitacca MD  
Mauro Carone MD  
Enrico Clini MD  
Mara Paneroni PT  
Marta Lazzeri PT  
Andrea Lanza PT  
Emilia Privitera PT  
Franco Pasqua MD  
Francesco Gigliotti MD  
Giorgio Castellana MD  
Paolo Banfi MD  
Enrico Guffanti MD  
Pierachille Santus MD  
Nicolino Ambrosino MD,

on behalf of the Italian Thoracic Society (ITS - AIPO), Association for the Rehabilitation of Respiratory Failure (ARIR) and the Italian Respiratory Society (SIP/IRS)

Graphics and publishing  
AIPO Ricerche Ed. - Milano  
edizioni@aiporicerche.it

Version - March 08, 2020

This document is available on  
<http://www.aiponet.it> and <https://www.arirassociazione.org/> and <http://www.sipirs.it/cms/>  
You can print all copies of this document for free.  
Copyright © 2020 by the AIPO – ITS, ARIR and SIP/IRS

## Introduction

The global outbreak of COVID-19 is having a dramatic effect throughout the world.<sup>1,2</sup> The widespread diffusion of the virus in Italy<sup>3</sup> has led national and regional authorities to institute extraordinary measures to contain the spread. The disease can cause major alveolar damage resulting in hypoxemic acute respiratory failure (ARF), which necessitates mechanical ventilation in a high proportion of cases.<sup>4-6</sup>

Italian societies of respiratory health care professionals promoted an audit among pulmonologists directly involved in the treatment of COVID-19 with the aim to formulate guidelines and propose a common management algorithm to apply in treating this condition in hospital setting.<sup>7</sup> The audit identified two urgent areas for action: organization and treatment. The latter, we believe, should also include respiratory rehabilitation, as it is an important component of the therapeutic management of these patients. Respiratory rehabilitation (RR) is defined as "*a multidisciplinary intervention based on personalized evaluation and treatment which includes, but is not limited to, exercise training, education, and behavioral modification designed to improve the physical and psychological condition of people with respiratory disease*".<sup>8</sup> In this context, and in relation to the topic of this document - patients affected by COVID-19 - physiotherapy interventions are particularly important.

Based on our general experience so far in dealing with the disease and on the existing knowledge (albeit still limited and evolving), we here propose a Consensus Statement on Respiratory Rehabilitation in patients with COVID-19. It represents an expert opinion of clinicians engaged in the field of RR, called - together with other specialties - to face with this emergency.

## Literature search

We carried out a search of the published literature in PubMed, Ovid, Embase online databases and relevant websites from the construction of databases to March 19, 2020 in order to retrieve guidelines and recommendations, meta-analyses, systematic reviews, state of the art, and randomized trials. The search terms were: "Coronavirus pneumonia", "ARDS", "Acute Hypoxemic Respiratory Failure", "SARS", "MERS", "Influenza", "Pulmonary Rehabilitation or Physical Therapy AND Infectious diseases, or Acute Respiratory Failure or Mechanical Ventilation" , "Noninvasive ventilation AND Acute hypoxemic respiratory failure", "Weaning from mechanical ventilation".

### **Generation of recommendations and consensus**

Based on the literature search, a small group of professionals produced a preliminary document which was then submitted to the participants. Consensus on the final document was achieved through a video conference meeting.

Below we summarize the recommendations that we consider most appropriate and urgent.

### ***General recommendations***

- The group takes into account the definition of severity based on Italian Health Organisation<sup>9</sup>:
  - a) mild condition: patient with pneumonia (clinical or radiological diagnosis) without any signs of severe pneumonia; b) severe condition: fever or suspected respiratory infection, respiratory frequency >30 b/min, dyspnoea, SpO<sub>2</sub> < 90% on room air; c) very severe condition (assumed to mimicking ARDS): bilateral interstitial radiological signs, paO<sub>2</sub>/FiO<sub>2</sub> <300

- Healthcare operators assigned to RR need to be sufficiently skilled (appropriate CV and experience with respiratory physiotherapy); if a dedicated staff is not present, health professionals must pass a training test on few but clear principles of behavior and action
- Operators and patients must follow all precautionary and preventive measures and wear all the protective gear specified in the relevant documents<sup>10-11</sup>
- RR is structured non-pharmacological therapy with a process delivered in three phases: assessment, intervention and re-assessment (the assessment is above all of a functional nature, with particular attention paid to the state of consciousness, respiratory, cardiological, motor function, and quality of life)
- All interventions and activities must be performed to avoid or reduce the risk of droplets production (particularly concerning airway clearance interventions)<sup>12</sup>
- Regardless of the type of intervention, the intensity, timing and modality must be tailored to the individual patient's needs, in particular for those with severe/critical illness, elderly patients, obesity, comorbidity and other complications<sup>13,14</sup>
- Assessment and monitoring should continue throughout the entire rehabilitation process. Patients who do not meet the standards set need to be reassessed daily.
- RR operators can also attend to reduce anxiety and depression in patients experiencing delirium, anger, fear, dysthymia, insomnia, panic attack, or sense of abandonment during isolation and intensive treatment, or exhibiting risk of non-collaboration or non-compliance with treatment.

**Acute phase (very severe respiratory condition)**

***Setting: Emergency Room, First aid, emergency care unit, intensive care unit (ICU), step-up sub-intensive care.***

- **The rehabilitative intervention** in this phase had to be started when patient has reached a minimum clinical stability. Treatments should be withdrawn in case of: high fever; worsening dyspnea; respiratory rate > 30 breaths/min; pulse oximetry <93% on oxygen therapy or requiring FiO<sub>2</sub> >50% during noninvasive ventilation (NIV), positive end expiratory pressure (PEEP)/continuous positive airway pressure (CPAP) >10 cm H<sub>2</sub>O; respiratory distress, arterial hypertension, brady- or tachycardia, intercurrent arrhythmias, shock, deep sedation, evidence of radiological lesions' progression (>50%) within 24-48 h<sup>15</sup>
- **The usual respiratory physiotherapy** procedures aimed at promoting the reduction of dyspnea, the tracheobronchial clearance, the training of the skeletal muscle and the maintenance / recovery of the ADL are contraindicated as they can determine a further loading of the respiratory system exposing the patient to an increased risk of distress
- When treating ARF by NIV or weaning from mechanical ventilation [MV] it is recommended to use similar protocols to those indicated for *de novo* ARF (e.g. FiO<sub>2</sub> reduction, PEEP reduction)<sup>16-19</sup>
- Airways clearance techniques are not recommended during the acute phase in patients without major problems of bronchial obstruction. Indeed, the hypothetical benefits do not outweigh the contamination risks for operators. The risk/benefit ratio should be evaluated on a single case basis in patients with bronchiectasis or with evident bronchial encumbrance, using tools that guarantee a safe distance from the patient<sup>15</sup>
- Twice per day evaluation of respiratory clinical parameters (temperature, SaO<sub>2</sub>, SpO<sub>2</sub>/FiO<sub>2</sub>, cough, dyspnea, respiratory rate, thoraco-abdominal dynamics) is indicated. We recommend to develop an algorithm to manage the assessment/intervention
- Evaluation of the peripheral muscle strength trend (by the Medical Research Council [MRC] scale and dynamometers) should be made as soon as practicable
- Frequent changes of posture, passive mobilization and/or neuromuscular electrical stimulation should be planned in the unconscious patient.<sup>15, 17-30</sup>

- Positional therapy (seated, semi-orthopneic, prone) with close monitoring is indicated in order to improve the ventilation/perfusion ratio and to prevent damage from immobilization.<sup>15, 17-30</sup>

**Table 1. Physiotherapy for COVID-19 patients in the critical phase**

<b>Ventilation support/Weaning</b>
○ Monitoring of clinical conditions (parameters and signs)
○ Adjustment of the mechanical support and oxygen therapy
○ Extubation protocols with or without NIV/CPAP
<b>Disability Prevention</b>
○ Passive/active mobilization
○ Frequent posture changes
○ Therapeutic postures (early sitting/pronation)
○ Neuromuscular electrical stimulation



## Acute phase (severe condition)

*Setting: Non-intensive acute care hospital (internal medicine, respiratory, infectious diseases, or other wards)*

- In hospitalized patients with mild/moderate disease, RR can improve symptoms (dyspnea, anxiety and depression), physical capacity and quality of life, at least potentially<sup>10-13, 17, 19-30</sup>
- Physiotherapy should be withdraw in case of: high fever, worsening dyspnea; pulse oximetry <93% or at least 4-point drop during exercise (desaturation), chest tightness, belching, dizziness, headache, unclear vision, palpitations, sweating, inability to keep balance, increased need for O<sub>2</sub> or NIV support, evidence of radiological lesions' progression (>50%) within 24-48 h<sup>15</sup>

**Table 2. Physiotherapy for COVID-19 patients in the acute phase**

<b>Ventilation support/Weaning</b>
○ Monitoring of clinical conditions (parameters and signs)
○ Adjustment of oxygen therapy
<b>Disability prevention</b>
○ Mobilization (getting patient out of bed)
○ Frequent posture changes/continuous rotational therapy
○ Therapeutic postures (early sitting/pronation)
○ Active limb exercises (also with dedicated devices) and muscle reconditioning
○ Strengthening the peripheral muscles
○ Neuromuscular electrical stimulation
○ Respiratory muscle training in case of inspiratory muscle weakness

## Chest physiotherapy

- Non-productive dry cough should be sedated to avoid fatigue and dyspnea
- Bronchial clearance techniques are indicated in hypersecretive patients with chronic respiratory diseases, by preferably using disposable devices with self-management (closed plastic bags for sputum collection help to prevent any spread of the virus).

## Post-Acute phase (post discharge from acute care to an intermediate setting)

**Setting:** *Respiratory rehabilitation department, other units for specialist and maintenance rehabilitation, intermediate care facilities, subacute wards*

- Similarly to patients recovering from ARDS due to H1N1 infection<sup>31-33</sup>, those with an acute COVID event may present disability and functional damage (respiratory function, critical illness myopathy and neuropathy), reduced participation and deterioration in quality of life, either in the short and long time following discharge
- Recovery time is variable depending upon the degree of normocapnic respiratory failure, and the associated physical (asthenia, peripheral muscle weakness) and emotional (anxiety, depression, sense of abandonment, post-traumatic stress syndrome) dysfunction<sup>34</sup>. Patients with comorbidities will take a longer period to return to their former condition
- Protocols for evaluating clinical parameters (temperature, SaO<sub>2</sub>, SpO<sub>2</sub>/FiO<sub>2</sub>, cough, dyspnea, respiratory rate, thoraco-abdominal dynamics) are indicated on a daily basis<sup>12,15</sup>
- Simple and repeatable treatment protocols to wean oxygen therapy are indicated
- It is recommended to evaluate peripheral muscle strength with the MRC scale, manual muscle test, isokinetic muscle test and measurement of joint range of motion (ROM)
- In weaned patients and those with prolonged weaning from mechanical ventilation and oxygen use, reconditioning interventions are indicated in order to improve the physical status

and to correct the motor and cognitive effects of prolonged immobilization in the intensive care area<sup>35-43</sup>

- As the effect of muscle activity in infections related to viral agents is not known, exercise aiming<sup>12</sup> with gradual load increase based on subjective symptoms are recommended to maintaining a normal function
- Low intensity exercise (<3.0 METs), daily patient counseling and education are indicated.<sup>12,15</sup>
- For patients i isolation, rehabilitation programs can be eventually conducted remotely by tele-health system (educational videos, tele-consultation, webcams etc., with disinfectable tools)<sup>12,15</sup>
- Patients discharged home or to other facilities in the community should receive indications on how to cope with physical activity, which need to be closely monitored regarding function, capacity and participation when the patient is cured with no longer risk of contagion
- An assessment of balance function is recommended as soon as possible (especially for patients who were bedridden for long)
- As soon as possible, assessment of exercise capacity, oxygenation response during effort (by the 6-min walk test-6MWT) and night-time should be planned.

**Table 3. Physiotherapy for COVID-19 patients during the post-acute phase**

<b>Weaning</b>
○ Monitoring of parameters
○ Mechanical ventilation weaning in tracheostomized patients
○ Management of problems related to tracheostomy (phonation, secretions encumbrance)

○ Adjustment of oxygen therapy
<b>Recovery from disability</b>
○ Mobilization (getting patient out of bed)
○ Frequent posture changes (sitting posture, prone positioning )
○ Strengthening of peripheral muscles
○ Reconditioning with specific aids (upper/lower limb devices/cycle-ergometer)
○ Neuromuscular electrical stimulation
○ Respiratory muscle training in case of inspiratory muscle weakness
<b>Chest physiotherapy</b>
○ Non-productive dry cough should be sedated to avoid fatigue and dyspnea
Bronchial clearance techniques are indicated in hypersecretive patients with chronic respiratory diseases, by preferably using disposable devices with self-management (closed plastic bags for sputum collection help to prevent any spread of the virus).
○ Pre-discharge counseling concerning physical activity
○ Support to the care team
■ ○ Psychological advice and support

■ [redacted]

■ [redacted]

### Organizational changes

The "war-time" emergency which developed within few weeks has forced the world of Rehabilitation to adapt to the circumstances and place itself at the disposal of the Regional/National Public Health Crisis Units.

- Hospitals dedicated to rehabilitation have expressed their willingness to set up, with flexibility, internal areas to cater for different categories of patients:

- a) isolated patients with suspected infection, awaiting confirmation of diagnosis;
  - b) symptomatic positive patients;
  - c) positive patients with respiratory failure on oxygen therapy, CPAP and NIV;
  - d) negative patients who require post COVID rehabilitation programs
- Rehabilitation teams coordinated by pulmonologists and respiratory therapists have now:
    - a) become part of the hospital programs for COVID-19 patients
    - b) made their services available to other rehabilitation professionals and professional nurses to create multidisciplinary groups able to support the high load of demand for care and assistance
    - c) shared with the other professionals protocols useful for clinical decision making "*ceiling of treatment*" (oxygen only, NIV/CPAP only, sedation), based on agreed criteria (patient characteristics, age, etc.)

### **Reorganization of the rehabilitation pathway for NON-COVID patients**

- Following the organizational changes determined by the COVID-19 epidemic, it is abundantly clear that all the other non infected patients with chronic respiratory diseases are suffering from these changes in their usual reference facilities as well as in the clinical and rehabilitative pathways
- For the full period of emergency and in the post-emergency period, the need for RR of patients not affected by COVID-19 infection will still have to be met, probably by new and diversified modalities of delivery (e.g. tele-rehabilitation paths, etc.).<sup>44,45</sup>

### **Conclusions**

The dramatic spread of the current COVID-19 epidemic in Italy has spurred into action also respiratory rehabilitation specialists (pulmonologists and respiratory therapists), who have been engaged for years in the care of patients with disabilities secondary to respiratory diseases and/or conditions. Their experience acquired in the management of chronic and acute respiratory failure is proving to be a fundamental asset for the management of patients during COVID-19 epidemic. Hence, it is likely that the reorganization involved in taking care of this scenario will not be a short-term matter.

### **Acknowledgments**

The Authors thank Laura Comini for editorial support and Rosemary Allpress for the English revision, and Giuseppe Gaudiello, Bruno Balbi, Ernesto Crisafulli, Mauro Maniscalco, Maria Aliani, Daniele Colombo, Giuseppe La Piana, Andrea Vianello for their clinical and scientific support.

## References

1. <https://www.worldometers.info/coronavirus/> (Last access on 19 March, 2020).
2. WHO. Global surveillance for human infection with coronavirus disease (COVID-2019). [https://www.who.int/publications-detail/global-surveillance-for-human-infection-with-novel-coronavirus-\(2019-ncov\)](https://www.who.int/publications-detail/global-surveillance-for-human-infection-with-novel-coronavirus-(2019-ncov)) (Last access on 19 March, 2020).
3. Istituto Superiore di Sanità. <https://www.epicentro.iss.it/coronavirus/>. Accessed on Centers for Disease Control and Prevention (CDC) 2019 Novel Coronavirus. <https://www.cdc.gov/> (Last access on 19 March, 2020).
4. Zhang Y, Xu J, Li H, Cao B. A novel coronavirus (COVID-19) outbreak: a call for action. *Chest*. 2020 Feb 19. pii: S0012-3692(20)30323-8. doi: 10.1016/j.chest.2020.02.014. [Epub ahead of print]
5. Zhu N, Zhang D, Wang W, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med*. 2020. *N Engl J Med*. 2020; 382: 727-733.
6. Huang C, Wang Y, Li X et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan China. *Lancet*. 2020;395:497-506.
7. Vitacca S, Nava S, Pintus A, Harari S. Facing the respiratory SARS-CoV-2 emergency in Italy: from ward to trenches. *Eur Respir J* 2020; in press.
8. Spruit MA, Singh SJ, Garvey C, et al. An official American Thoracic Society/European Respiratory Society statement: key concepts and advances in pulmonary rehabilitation. *Am J Respir Crit Care Med*. 2013; 188: e13-64.
9. Istituto Nazionale per le Malattie Infettive Lazzaro Spallanzani – IRCCS - Protocollo di gestione clinica e raccolta dati dei casi confermati di Malattia da Infezione da Coronavirus – 2019 (COVID-19) nel paziente adulto- 19 March 2020 - A cura dell'INMI COVID-19 group, con la collaborazione della Rete Regionale di Malattie infettive, dei PS/DEA e delle Terapie Intensive. <https://www.inmi.it/seresmi>. (Last access 26 March 2020).
10. [https://apps.who.int/iris/bitstream/handle/10665/331215/WHO-2019-nCov-IPCPPE\\_use-2020.1-eng.pdf](https://apps.who.int/iris/bitstream/handle/10665/331215/WHO-2019-nCov-IPCPPE_use-2020.1-eng.pdf). (Last access 19 March 2020).
11. Circolare Ministero della Salute n. 5443 del 22 February 2020 [http://www.prefettura.it/FILES/allegatinews/1181/Circolare\\_Ministero\\_della\\_Salute\\_n.\\_5443\\_del\\_22\\_febbraio\\_2020.pdf](http://www.prefettura.it/FILES/allegatinews/1181/Circolare_Ministero_della_Salute_n._5443_del_22_febbraio_2020.pdf). (Last access 24 Marzo 2020).
12. Zhonghua Jie He He Hu Xi Za Zhi. Recommendations for respiratory rehabilitation of COVID-19 in adult. 2020 Mar 3;43(0):E029. doi: 10.3760/cma.j.cn112147-20200228-00206. [Epub ahead of print]
13. Ambrosino N, Clini EM. Response to pulmonary rehabilitation: toward personalised programmes? *Eur Respir J*. 2015; 46 (6): 1538-40.
14. Wouters EFM, Wouters BBREF, Augustin IML, Houben-Wilke S, Vanfleteren LEGW, Franssen FME. Personalised pulmonary rehabilitation in COPD. *Eur Respir Rev*. 2018;27:170125.
15. The first affiliated Hospital, Zhejiang University School of Medicine. Compiled According to Clinical Experience. Rehabilitation therapy for COVID-19 patients. In: Handbook of COVID-19, prevention and treatment 2020: pp 47-48. <http://www.zju.edu.cn/english/2020/0323/c19573a1987520/page.htm>. (Last Access 27 March 2020).
16. Esquinas AM, Pravinkumar SE, Scala R, et al. Noninvasive mechanical ventilation in high-risk pulmonary infections: a clinical review. *Eur Respir Rev*. 2014; 23 (134): 427-438.
17. Gosselink R, Clini E. Rehabilitation in Intensive Care. In: Clini E, Holland A, Pitta F, Troosters T. (eds) *Textbook of Pulmonary Rehabilitation*. Springer Nature, Cham (CH), 2018: pp 349-366.
18. van der Lee L, Hill AM, Patman S. Expert consensus for respiratory physiotherapy management of mechanically ventilated adults with community-acquired pneumonia: A Delphi study. *J Eval Clin Pract*. 2019; 25 (2):230–243.
19. Doiron KA, Hoffmann TC, Beller EM. Early intervention (mobilization or active exercise) for critically ill adults in the intensive care unit (Review). *Cochrane Database System Rev*. 2018; 3: CD010754.
20. Ambrosino N, Makhbah DN. Comprehensive physiotherapy management in ARDS. *Minerva Anestesiol*. 2013; 79 (5):554-563.

21. Schweickert WD, Pohlman MC, Pohlman AS et al. Early physical and occupational therapy in mechanically ventilated, critically ill patients: A randomised controlled trial. *Lancet*. 2009; 373 (9678):1874–1882.
22. Connolly B, O'Neill B, Salisbury L, Blackwood B. Physical rehabilitation interventions for adult patients during critical illness: An overview of systematic reviews. *Thorax*. 2016; 71(10):881–90.
23. Hanekom S, Gosselink R, Dean E et al. The development of a clinical management algorithm for early physical activity and mobilization of critically ill patients: Synthesis of evidence and expert opinion and its translation into practice. *Clin Rehabil*. 2011; 25 (9):771–787.
24. Segers J, Hermans G, Bruyninckx F, et al. Feasibility of neuromuscular electrical stimulation in critically ill patients. *J Crit Care*. 2014; 29 (6):1082–1088.
25. Guarracino F, Bertini P, Bortolotti U, et al. Flexible bronchoscopy during mechanical ventilation in the prone position to treat acute lung injury. *Rev Port Pneumol*. 2013; 19 (1): 42-44.
26. Medrinal C, Combret Y, Prieur G et al. Comparison of exercise intensity during four early rehabilitation techniques in sedated and ventilated patients in ICU: A randomised cross-over trial. *Crit Care*. 2018; 22 (1):110.
27. Gattinoni L, Tognoni G, Pesenti A et al.; Prone-Supine Study Group. Effect of prone positioning on the survival of patients with acute respiratory failure. *N Engl J Med*. 2001; 345 (8): 568–573.
28. Karatzanos E, Gerovasili V, Zervakis D et al. Electrical muscle stimulation: An effective form of exercise and early mobilization to preserve muscle strength in critically ill patients. *Crit Care Res Pract*. 2012; 2012: 432752.
29. Fossat G, Baudin F, Courtes L et al. Effect of in-bed leg cycling and electrical stimulation of the quadriceps on global muscle strength in critically ill adults: A Randomized Clinical Trial. *JAMA*. 2018; 320 (4): 368–378.
30. Fuke R, Hifumi T, Kondo Y et al. Early rehabilitation to prevent postintensive care syndrome in patients with critical illness: A systematic review and meta-analysis. *BMJ Open*. 2018; 8 (5): e019998.
31. Hsieh, Meng - Jer, et al. Recovery of pulmonary functions, exercise capacity, and quality of life after pulmonary rehabilitation in survivors of ARDS due to severe influenza A (H1N1) pneumonitis. *Influenza Other Respir Viruses*. 2018; 12 (5): 643-648.
32. Orme J, Romney JS, Hopkins RO, et al. Pulmonary Function and Health-related Quality of Life in Survivors of Acute Respiratory Distress Syndrome. *Am J Respir Crit Care Med* 2003; 167 (5): 690-94.
33. Hill DA, Fowler RA, Burns KE, et al. Long-term outcomes and healthcare utilization after prolonged mechanical ventilation. *Ann Am Thorac*. 2017; 14 (3): 355–362.
34. Pandharipande PP, Girard TD, Jackson JC et al.; BRAIN-ICU Study Investigators. Long-term cognitive impairment after critical illness. *N Engl J Med*. 2013; 369 (14):1306–1316.
35. Girard TD, Alhazzani W, Kress JP et al.; ATS/CHEST Ad Hoc Committee on Liberation from Mechanical Ventilation in Adults. An official American Thoracic Society/American College of Chest Physicians Clinical Practice Guideline: Liberation from mechanical ventilation in critically ill adults. Rehabilitation protocols, ventilator liberation protocols, and cuff leak tests. *Am J Respir Crit Care Med*. 2017; 195 (1): 120–133.
36. Vitacca M, Clini EM, Nava S, Ambrosino N. La Riabilitazione ad Alta Complessità nel Paziente con Svezzamento Prolungato: Ruolo dello Pneumologo. *Rass Patol App Resp* 2013; 28: 179-187. [Italian].
37. Ambrosino N, Vitacca M. The patient needing prolonged mechanical ventilation: A narrative review. *Multidiscip Respir Med*. 2018; 13: 6.
38. Ceriana P, Nava S, Vitacca M, et al. Noninvasive ventilation during weaning from prolonged mechanical ventilation. *Pulmonology*. 2019; 25 (6):328-333.
39. Winck JC, Gilet H, Kalin P, et al. Validation of the Multi-INdependence Dimensions (MIND) questionnaire for prolonged mechanically ventilated subjects. *BMC Pulm Med*. 2019; 19 (1):109.
40. Schreiber AF, Ceriana P, Ambrosino N, et al. Physiotherapy and weaning from prolonged mechanical ventilation. *Respir Care*. 2019; 64 (1):17-25.
41. Ambrosino N, Venturelli E, Vaghegini G, Clini E. Rehabilitation, weaning and physical therapy strategies in chronic critically ill patients. *Eur Respir J*. 2012; 39 (2): 487-92.
42. Bissett BM, Leditschke IA, Neeman T, et al. Inspiratory muscle training to enhance recovery from mechanical ventilation: A randomised trial. *Thorax*. 2016; 71 (9): 812–819.
43. Vorona S, Sabatini U, Al-Maqbali S et al. Inspiratory muscle rehabilitation in critically ill adults. A systematic review and meta-analysis. *Ann Am Thorac Soc*. 2018; 15 (6): 735–744.



44. Donner CF, Raskin J, ZuWallack R, et al. Incorporating telemedicine into the integrated care of the COPD patient a summary of an interdisciplinary workshop held in Stresa, Italy, 7-8 September 2017. *Respir Med.* 2018; 143: 91-102.
45. Ambrosino N, Fracchia C. The role of tele-medicine in patients with respiratory diseases. *Expert Rev Respir Med.* 2017;11 (11): 893-900.